

PART 7

Supplementary tables

SUPPLEMENTARY TABLES

List of tables

Distribution of the Young Minds Matter survey sample 189

Table S-1: Parent or carer interview sample by sex and age group	189
Table S-2: Parent or carer interview sample by family type	189
Table S-3: Parent or carer interview sample by household income	189
Table S-4: Parent or carer interview sample by parent or carer education level	190
Table S-5: Parent or carer interview sample by parent or carer labour force status	190
Table S-6: Parent or carer interview sample by area of residence	190
Table S-7: Parent or carer interview sample by country of birth of child or adolescent.....	190
Table S-8: Parent or carer interview sample by country of birth of parents or carers	191
Table S-9: Parent or carer interview sample by housing tenure.....	191
Table S-10: Parent or carer interview sample by remoteness area.....	191
Table S-11: Parent or carer interview sample by index of relative socio-economic disadvantage.....	191
Table S-12: 4-17 year-olds by level of family functioning.....	192

Prevalence of mental disorders..... 192

Table S-13: 12-month prevalence of mental disorder among 4-17 year-olds by country of birth of child or adolescent.....	192
Table S-14: 12-month prevalence of mental disorder among 4-17 year-olds by country of birth of parents or carers	192
Table S-15: 12-month prevalence of mental disorder among 4-17 year-olds by housing tenure.....	192
Table S-16: 12-month prevalence of mental disorder among 4-17 year-olds by remoteness area	193
Table S-17: 12-month prevalence of mental disorder among 4-17 year-olds by index of relative socio-economic disadvantage.....	193

Principal condition..... 193

Table S-18: Principal condition by sex and age group	193
--	-----

Major depressive disorder 194

Table S-19: 12-month prevalence of major depressive disorder among 4-17 year-olds by country of birth of child or adolescent	194
Table S-20: 12-month prevalence of major depressive disorder among 4-17 year-olds by country of birth of parents or carers.....	194
Table S-21: 12-month prevalence of major depressive disorder among 4-17 year-olds by housing tenure.....	194
Table S-22: 12-month prevalence of major depressive disorder among 4-17 year-olds by remoteness area..	194

Table S-23: 12-month prevalence of major depressive disorder among 4-17 year-olds by index of relative socio-economic disadvantage.....	195
Anxiety disorders	195
Table S-24: 12-month prevalence of anxiety disorders among 4-17 year-olds by country of birth of child or adolescent.....	195
Table S-25: 12-month prevalence of anxiety disorders among 4-17 year-olds by country of birth of parents or carers	195
Table S-26: 12-month prevalence of anxiety disorders among 4-17 year-olds by housing tenure.....	196
Table S-27: 12-month prevalence of anxiety disorders among 4-17 year-olds by remoteness area	196
Table S-28: 12-month prevalence of anxiety disorders among 4-17 year-olds by index of relative socio-economic disadvantage.....	196
ADHD	197
Table S-29: 12-month prevalence of ADHD among 4-17 year-olds by country of birth of child or adolescent	197
Table S-30: 12-month prevalence of ADHD among 4-17 year-olds by country of birth of parents or carers.....	197
Table S-31: 12-month prevalence of ADHD among 4-17 year-olds by housing tenure	197
Table S-32: 12-month prevalence of ADHD among 4-17 year-olds by remoteness area	197
Table S-33: 12-month prevalence of ADHD among 4-17 year-olds by index of relative socio-economic disadvantage.....	198
Conduct disorder	198
Table S-34: 12-month prevalence of conduct disorder among 4-17 year-olds by country of birth of child or adolescent.....	198
Table S-35: 12-month prevalence of conduct disorder among 4-17 year-olds by country of birth of parents or carers	198
Table S-36: 12-month prevalence of conduct disorder among 4-17 year-olds by housing tenure.....	198
Table S-37: 12-month prevalence of conduct disorder among 4-17 year-olds by remoteness area.....	199
Table S-38: 12-month prevalence of conduct disorder among 4-17 year-olds by index of relative socio-economic disadvantage.....	199
Mental disorders in adolescents aged 16-17 years.....	199
Table S-39: 12-month prevalence of mental disorders among 16-17 year-olds by type of disorder and sex ...	199
Table S-40: 12-month prevalence of mental disorders among 16-17 year-olds by type of disorder and severity of impact.....	200
Table S-41: 12-month prevalence of oppositional problem behaviours among 16-17 year-olds by sex.....	200
Table S-42: Average days absent from school in past 12 months due to symptoms of mental disorders among 16-17 year-olds with mental disorders by type of disorder	200
Strengths and Difficulties Questionnaire	201
Table S-43: Prevalence of parent or carer reported Strengths and Difficulties Questionnaire (SDQ) total difficulties score in the abnormal range among 4-17 year-olds by sex and age group.....	201

Table S-44: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale, sex and age group	201
Table S-45: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and household income	201
Table S-46: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and highest level of education of either primary or secondary parent or carer	202
Table S-47: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and labour force status of parents or carers	202
Table S-48: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and area of residence	202
Table S-49: 4-17 year-olds by parent or carer reported SDQ total difficulties score and SDQ impact score	203
K10+ days out of role.....	203
Table S-50: Number of days out of role (K10+) among 11-17 year-olds by sex.....	203
Self-harm	204
Table S-51: Self-harm among 12-17 year-olds by sex and adolescent self-reported SDQ rating	204
Table S-52: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and family type	205
Table S-53: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and household income	206
Table S-54: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer education.....	206
Table S-55: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer labour force status	207
Table S-56: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and area of residence	208
Suicidal behaviours.....	208
Table S-57: Suicide ideation, suicide plans and suicide attempts among 12-17 year-olds by sex and adolescent self-reported SDQ rating	208
Table S-58: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and family type	209
Table S-59: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and household income	210
Table S-60: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer education.....	210
Table S-61: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer labour force status	211
Table S-62: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and area of residence	211

Smoking, alcohol, and cannabis use and sexual behaviours by Strengths and Difficulties**Questionnaire scales..... 212**

Table S-63: Smoking among 13-17 year-olds by sex and adolescent self-reported SDQ rating.....212

Table S-64: Alcohol consumption among 13-17 year-olds by sex and adolescent self-reported SDQ rating...213

Table S-65: Cannabis and other drug use among 13-17 year-olds by sex and adolescent self-reported SDQ rating214

Table S-66: Sexual behaviours among 13-17 year-olds by sex and adolescent self-reported SDQ rating.....215

Table S-67: Use of protection during last sexual intercourse or consumption of alcohol or drugs prior to last sexual intercourse among 13-17 year-olds who have ever had sexual intercourse by sex and adolescent self-reported SDQ rating.....216

Problem eating behaviours 217

Table S-68: Problem eating behaviours and weight control behaviours in the past 12 months among 11-17 year-olds by sex and age group217

Bullying..... 219

Table S-69: Frequency of bullying in previous 12 months in 11-17 year-olds by sex and age group.....219

Internet use and electronic games 220

Table S-70: Time spent using the internet or playing electronic games among 11-17 year-olds by sex and age group220

Table S-71: Problem internet or electronic gaming behaviours among 11-17 year-olds by sex and age group221

Table S-72: Problem internet use or electronic gaming behaviours among 11-17 year-olds by sex and adolescent self-reported SDQ rating.....221

Distribution of the Young Minds Matter survey sample

Table S-1: Parent or carer interview sample by sex and age group

Sex	Age group	Respondents	Proportion (%)
Males	4-11 years	1709	27.1
	12-17 years	1545	24.5
	4-17 years	3254	51.6
Females	4-11 years	1625	25.8
	12-17 years	1431	22.7
	4-17 years	3056	48.4
Persons	4-11 years	3334	52.8
	12-17 years	2976	47.2
	4-17 years	6310	100.0

Table S-2: Parent or carer interview sample by family type

Family type	Respondents	Proportion (%)
Families with two parents or carers	5060	80.2
Original family	4331	68.6
Step family	283	4.5
Blended family	392	6.2
Other family	54	0.9
Families with one parent or carer	1250	19.8
Total	6310	100.0

'Original family' has at least one child living with their natural, adoptive or foster parents, and no step children.

'Other family' is where all children are not the natural, adopted, foster or step child of one or both carers.

Table S-3: Parent or carer interview sample by household income

Household income before tax	Respondents	Proportion (%)
\$130,000 or more per year	1686	26.7
\$52,000-\$129,999 per year	2833	44.9
Less than \$52,000 per year	1479	23.4
Not stated	312	4.9
Total	6310	100.0

Household income includes the combined income for the 2011-12 financial year of everyone living in the household before tax and other deductions are taken out.

Table S-4: Parent or carer interview sample by parent or carer education level

Highest level of education of either primary or secondary parent or carer	Respondents	Proportion (%)
Bachelor degree or higher	2667	42.3
Diploma or certificate III/IV	2414	38.3
Year 11 or 12	756	12.0
Year 10 or below	473	7.5
Total	6310	100.0

Table S-5: Parent or carer interview sample by parent or carer labour force status

Parent or carer labour force status	Respondents	Proportion (%)
Both parents or carers employed	3413	54.1
One parent or carer employed, one parent or carer not in employment	1455	23.1
Both parents or carers not in employment	209	3.3
Sole parent or carer employed	740	11.7
Sole parent or carer not in employment	475	7.5
Not stated	18	0.3
Total	6310	100.0

'Not in employment' combines unemployed and not in the labour force.

Table S-6: Parent or carer interview sample by area of residence

Area of residence	Respondents	Proportion (%)
Greater capital cities	4044	64.1
Rest of state	2266	35.9
Total	6310	100.0

Based on the ABS classification Greater Capital City Statistical Area (GCCSA).

Table S-7: Parent or carer interview sample by country of birth of child or adolescent

Child or adolescent country of birth	Respondents	Proportion (%)
Australia	5540	87.8
Overseas	770	12.2
Total	6310	100.0

Table S-8: Parent or carer interview sample by country of birth of parents or carers

Parent or carer country of birth	Respondents	Proportion (%)
Both parents or carers born in Australia	2977	47.2
One parent or carer born in Australia, one parent or carer born overseas	929	14.7
Both parents or carers born overseas	1188	18.8
Sole parent or carer born in Australia	956	15.2
Sole parent or carer born overseas	259	4.1
Not stated	1	0.0
Total	6310	100.0

Table S-9: Parent or carer interview sample by housing tenure

Housing tenure	Respondents	Proportion (%)
Owned outright	777	12.3
Owned with a mortgage	3488	55.3
Rented — public housing	268	4.2
Rented — other	1677	26.6
Other	88	1.4
Not stated	12	0.2
Total	6310	100.0

Table S-10: Parent or carer interview sample by remoteness area

Remoteness area	Respondents	Proportion (%)
Major Cities of Australia	4134	65.5
Inner Regional Australia	1554	24.6
Outer Regional Australia	516	8.2
Remote Australia or Very Remote Australia	106	1.7
Total	6310	100.0

Table S-11: Parent or carer interview sample by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Respondents	Proportion (%)
Lowest quintile (most disadvantaged)	1055	16.7
Second quintile	1180	18.7
Third quintile	1188	18.8
Fourth quintile	1436	22.8
Highest quintile (least disadvantaged)	1451	23.0
Total	6310	100.0

Table S-12: 4-17 year-olds by level of family functioning

Level of family functioning	Proportion (%)
Very good	60.8
Good	23.0
Fair	12.5
Poor	3.7

Family functioning was measured across a number of domains by the McMaster Family Assessment Device (see glossary).

Prevalence of mental disorders

Table S-13: 12-month prevalence of mental disorder among 4-17 year-olds by country of birth of child or adolescent

Child or adolescent country of birth	Prevalence (%)
Australia	14.8
Overseas	7.2

Table S-14: 12-month prevalence of mental disorder among 4-17 year-olds by country of birth of parents or carers

Parent or carer country of birth	Prevalence (%)
Both carers born in Australia	14.3
One carer born in Australia, one carer born overseas	12.9
Both carers born overseas	5.1
Sole carer born in Australia	24.6
Sole carer born overseas	13.4

Table S-15: 12-month prevalence of mental disorder among 4-17 year-olds by housing tenure

Housing tenure	Prevalence (%)
Owned outright	10.5
Owned with a mortgage	11.2
Rented — public housing	29.2
Rented — other	17.7
Other	18.2

Table S-16: 12-month prevalence of mental disorder among 4-17 year-olds by remoteness area

Remoteness area	Prevalence (%)
Major Cities of Australia	12.9
Inner Regional Australia	14.8
Outer Regional Australia	19.0
Remote Australia or Very Remote Australia	14.0

Table S-17: 12-month prevalence of mental disorder among 4-17 year-olds by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Prevalence (%)
Lowest quintile (most disadvantaged)	20.7
Second quintile	14.4
Third quintile	13.3
Fourth quintile	11.7
Highest quintile (least disadvantaged)	10.9

Principal condition

Table S-18: Principal condition by sex and age group

Disorder	Males 4-11 years (%)	Males 12-17 years (%)	Males 4-17 years (%)	Females 4-11 years (%)	Females 12-17 years (%)	Females 4-17 years (%)	Persons 4-11 years (%)	Persons 12-17 years (%)	Persons 4-17 years (%)
Social phobia	0.9	2.1	1.4	0.8	1.7	1.2	0.8	1.9	1.3
Separation anxiety	4.0	1.6	3.0	3.5	1.7	2.7	3.8	1.6	2.9
Generalised anxiety	1.1	0.8	1.0	0.7	1.6	1.1	0.9	1.2	1.0
Obsessive- compulsive disorder	np	np	0.2	np	np	0.1	0.1	0.2	0.2
Major depressive disorder	0.5	3.0	1.6	0.6	4.7	2.4	0.6	3.8	2.0
Conduct disorder	1.1	1.2	1.1	1.0	1.0	1.0	1.0	1.1	1.1
ADHD	8.6	6.9	7.9	4.0	1.8	3.1	6.3	4.5	5.5

np Not available for publication because of small cell size, but included in totals where applicable.

Major depressive disorder

Table S-19: 12-month prevalence of major depressive disorder among 4-17 year-olds by country of birth of child or adolescent

Child or adolescent country of birth	Prevalence (%)
Australia	3.0
Overseas	1.3

Table S-20: 12-month prevalence of major depressive disorder among 4-17 year-olds by country of birth of parents or carers

Parent or carer country of birth	Prevalence (%)
Both carers born in Australia	2.3
One carer born in Australia, one carer born overseas	2.8
Both carers born overseas	1.0
Sole carer born in Australia	6.2
Sole carer born overseas	3.0

Table S-21: 12-month prevalence of major depressive disorder among 4-17 year-olds by housing tenure

Housing tenure	Prevalence (%)
Owned outright	2.7
Owned with a mortgage	2.1
Rented — public housing	6.3
Rented — other	3.5
Other	4.4

Table S-22: 12-month prevalence of major depressive disorder among 4-17 year-olds by remoteness area

Remoteness area	Prevalence (%)
Major Cities of Australia	2.7
Inner Regional Australia	2.8
Outer Regional Australia	3.3
Remote Australia or Very Remote Australia	4.4

Table S-23: 12-month prevalence of major depressive disorder among 4-17 year-olds by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Prevalence (%)
Lowest quintile (most disadvantaged)	4.0
Second quintile	2.6
Third quintile	2.9
Fourth quintile	2.1
Highest quintile (least disadvantaged)	2.6

Anxiety disorders

Table S-24: 12-month prevalence of anxiety disorders among 4-17 year-olds by country of birth of child or adolescent

Child or adolescent country of birth	Social phobia (%)	Separation anxiety (%)	Generalised anxiety (%)	Obsessive-compulsive (%)	Any anxiety disorder (%)
Australia	2.5	4.6	2.4	0.8	7.4
Overseas	1.1	1.4	np	0.9	3.2

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-25: 12-month prevalence of anxiety disorders among 4-17 year-olds by country of birth of parents or carers

Parent or carer country of birth	Social phobia (%)	Separation anxiety (%)	Generalised anxiety (%)	Obsessive-compulsive (%)	Any anxiety disorder (%)
Both carers born in Australia	2.1	4.0	2.3	0.7	6.6
One carer born in Australia, one carer born overseas	2.4	3.0	2.1	0.4	6.1
Both carers born overseas	0.7	0.9	0.5	0.5	2.1
Sole carer born in Australia	4.9	10.3	4.2	2.0	14.8
Sole carer born overseas	2.5	4.5	np	np	5.7

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-26: 12-month prevalence of anxiety disorders among 4-17 year-olds by housing tenure

Housing tenure	Social phobia (%)	Separation anxiety (%)	Generalised anxiety (%)	Obsessive-compulsive (%)	Any anxiety disorder (%)
Owned outright	1.8	1.7	1.3	0.6	4.1
Owned with a mortgage	1.8	3.4	1.9	0.5	5.8
Rented — public housing	6.0	12.1	4.3	3.5	16.6
Rented — other	2.9	5.5	2.7	1.1	8.5
Other	np	6.5	np	np	8.4

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-27: 12-month prevalence of anxiety disorders among 4-17 year-olds by remoteness area

Remoteness area	Social phobia (%)	Separation anxiety (%)	Generalised anxiety (%)	Obsessive-compulsive (%)	Any anxiety disorder (%)
Major Cities of Australia	2.2	3.9	2.1	0.9	6.4
Inner Regional Australia	3.0	5.2	2.5	0.9	8.1
Outer Regional Australia	1.7	4.4	1.6	np	7.7
Remote Australia or Very Remote Australia	np	3.9	np	np	5.3

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-28: 12-month prevalence of anxiety disorders among 4-17 year-olds by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Social phobia (%)	Separation anxiety (%)	Generalised anxiety (%)	Obsessive-compulsive (%)	Any anxiety disorder (%)
Lowest quintile (most disadvantaged)	3.8	6.1	2.5	2.0	10.4
Second quintile	1.6	5.1	2.2	0.7	7.3
Third quintile	2.9	3.6	2.2	0.4	6.2
Fourth quintile	1.6	3.6	2.3	0.4	6.1
Highest quintile (least disadvantaged)	2.0	3.2	1.7	0.8	5.3

ADHD**Table S-29: 12-month prevalence of ADHD among 4-17 year-olds by country of birth of child or adolescent**

Child or adolescent country of birth	Prevalence (%)
Australia	7.9
Overseas	3.9

Table S-30: 12-month prevalence of ADHD among 4-17 year-olds by country of birth of parents or carers

Parent or carer country of birth	Prevalence (%)
Both carers born in Australia	7.9
One carer born in Australia, one carer born overseas	6.9
Both carers born overseas	3.0
Sole carer born in Australia	12.1
Sole carer born overseas	6.4

Table S-31: 12-month prevalence of ADHD among 4-17 year-olds by housing tenure

Housing tenure	Prevalence (%)
Owned outright	5.3
Owned with a mortgage	5.8
Rented — public housing	17.2
Rented — other	9.6
Other	9.5

Table S-32: 12-month prevalence of ADHD among 4-17 year-olds by remoteness area

Remoteness area	Prevalence (%)
Major Cities of Australia	7.0
Inner Regional Australia	7.8
Outer Regional Australia	10.0
Remote Australia or Very Remote Australia	4.3

Table S-33: 12-month prevalence of ADHD among 4-17 year-olds by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Prevalence (%)
Lowest quintile (most disadvantaged)	12.2
Second quintile	8.1
Third quintile	7.0
Fourth quintile	5.6
Highest quintile (least disadvantaged)	5.2

Conduct disorder

Table S-34: 12-month prevalence of conduct disorder among 4-17 year-olds by country of birth of child or adolescent

Child or adolescent country of birth	Prevalence (%)
Australia	2.3
Overseas	0.5

Table S-35: 12-month prevalence of conduct disorder among 4-17 year-olds by country of birth of parents or carers

Parent or carer country of birth	Prevalence (%)
Both carers born in Australia	2.0
One carer born in Australia, one carer born overseas	1.0
Both carers born overseas	np
Sole carer born in Australia	5.2
Sole carer born overseas	4.3

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-36: 12-month prevalence of conduct disorder among 4-17 year-olds by housing tenure

Housing tenure	Prevalence (%)
Owned outright	0.6
Owned with a mortgage	1.2
Rented — public housing	10.8
Rented — other	3.0
Other	np

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-37: 12-month prevalence of conduct disorder among 4-17 year-olds by remoteness area

Remoteness area	Prevalence (%)
Major Cities of Australia	1.6
Inner Regional Australia	3.0
Outer Regional Australia	2.6
Remote Australia or Very Remote Australia	np

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-38: 12-month prevalence of conduct disorder among 4-17 year-olds by index of relative socio-economic disadvantage

Index of relative socio-economic disadvantage	Prevalence (%)
Lowest quintile (most disadvantaged)	5.7
Second quintile	1.4
Third quintile	1.6
Fourth quintile	1.1
Highest quintile (least disadvantaged)	1.1

Mental disorders in adolescents aged 16-17 years

Table S-39: 12-month prevalence of mental disorders among 16-17 year-olds by type of disorder and sex

Disorder	Males (%)	Females (%)	Persons (%)
Social phobia	2.8	4.8	3.8
Separation anxiety disorder	2.2	4.3	3.2
Generalised anxiety disorder	2.5	5.3	3.9
Obsessive-compulsive disorder	np	1.2	0.8
Any anxiety disorder	5.4	10.2	7.8
Major depressive disorder	5.4	10.0	7.7
ADHD	5.3	2.8	4.0
Conduct disorder	2.5	1.2	1.8
Any mental disorder	14.1	15.3	14.7

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-40: 12-month prevalence of mental disorders among 16-17 year-olds by type of disorder and severity of impact

Disorder	Mild (%)	Moderate (%)	Severe (%)	Total (%)
Social phobia	1.1	1.0	1.7	3.8
Separation anxiety disorder	0.8	1.0	1.4	3.2
Generalised anxiety disorder	1.1	1.5	1.3	3.9
Obsessive-compulsive disorder	np	np	0.3	0.8
Any anxiety disorder	2.6	2.6	2.7	7.8
Major depressive disorder	1.5	3.1	3.1	7.7
ADHD	2.2	1.4	0.4	4.0
Conduct disorder	0.6	0.9	0.3	1.8
Any mental disorder	5.2	5.8	3.7	14.7

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-41: 12-month prevalence of oppositional problem behaviours among 16-17 year-olds by sex

Sex	Per cent
Males	4.5
Females	6.6
Persons	5.5

Table S-42: Average days absent from school in past 12 months due to symptoms of mental disorders among 16-17 year-olds with mental disorders by type of disorder

Disorder	Average (days)
Social phobia	19
Separation anxiety disorder	41
Generalised anxiety disorder	19
Obsessive-compulsive disorder	15
Any anxiety disorder	24
Major depressive disorder	26
ADHD	8
Conduct disorder	7

Strengths and Difficulties Questionnaire

Table S-43: Prevalence of parent or carer reported Strengths and Difficulties Questionnaire (SDQ) total difficulties score in the abnormal range among 4-17 year-olds by sex and age group

Sex	Age group	Prevalence (%)	Population estimate
Males	4-11 years	11.8	140,000
	12-17 years	12.3	107,000
	4-17 years	12.0	247,000
Females	4-11 years	7.8	88,200
	12-17 years	8.3	68,900
	4-17 years	8.0	157,000
Persons	4-11 years	9.9	228,000
	12-17 years	10.3	176,000
	4-17 years	10.1	404,000

Table S-44: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale, sex and age group

SDQ scale	Males 4-11 years (%)	Males 12-17 years (%)	Females 4-11 years (%)	Females 12-17 years (%)	Persons 4-17 years (%)
Emotional problems	10.2	13.7	11.8	19.3	13.3
Conduct problems	12.5	11.6	9.5	7.9	10.5
Hyperactivity	19.0	14.9	10.2	6.4	13.0
Peer problems	12.6	16.4	9.6	13.4	12.7
Prosocial behaviour	2.9	4.5	1.9	1.8	2.7
Total difficulties	11.8	12.3	7.8	8.3	10.1

Table S-45: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and household income

Household income before tax	Emotional problems	Conduct problems	Hyperactivity	Peer problems	Prosocial behaviour	Total difficulties
\$130,000 or more per year	10.4	7.2	10.8	8.4	2.2	6.4
\$52,000-\$129,999 per year	13.0	9.7	11.4	11.6	2.3	9.0
Less than \$52,000 per year	16.5	15.2	18.2	18.5	4.0	15.2

Household income includes the combined income for the 2011-12 financial year of everyone living in the household before tax and other deductions are taken out.

Table S-46: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and highest level of education of either primary or secondary parent or carer

Highest level of education of either primary or secondary parent or carer	Emotional problems	Conduct problems	Hyperactivity	Peer problems	Prosocial behaviour	Total difficulties
Bachelor degree or higher	10.1	6.7	10.2	9.2	2.4	6.1
Diploma or certificate III/IV	14.9	12.5	14.2	14.0	2.7	11.8
Year 11 or 12	15.2	13.3	14.8	15.7	2.4	12.2
Year 10 or below	19.0	16.4	19.4	20.1	5.0	18.6

Table S-47: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and labour force status of parents or carers

Parent or carer labour force status	Emotional problems	Conduct problems	Hyperactivity	Peer problems	Prosocial behaviour	Total difficulties
Both parents or carers employed	11.2	7.7	10.6	10.2	2.3	7.4
One parent or carer employed, one parent or carer not in employment	13.4	11.6	13.3	12.9	2.3	10.6
Both parents or carers not in employment	17.0	19.6	18.8	17.5	4.9	15.9
Sole parent or carer employed	16.5	11.3	13.5	13.1	3.3	10.8
Sole parent or carer not in employment	20.8	19.8	24.5	25.5	5.2	22.0

Table S-48: Prevalence of parent or carer reported SDQ scale scores in the abnormal range among 4-17 year-olds by SDQ scale and area of residence

Area of residence	Emotional problems	Conduct problems	Hyperactivity	Peer problems	Prosocial behaviour	Total difficulties
Greater capital cities	13.1	9.5	12.7	11.0	2.5	9.5
Rest of state	13.7	12.2	13.7	15.6	3.1	11.1

Based on the ABS classification Greater Capital City Statistical Area (GCCSA).

Table S-49: 4-17 year-olds by parent or carer reported SDQ total difficulties score and SDQ impact score

Total Difficulties	SDQ impact – Normal	SDQ impact – Borderline	SDQ impact – Abnormal
Normal	89.4	4.7	5.9
Borderline	39.6	17.5	42.9
Abnormal	16.7	11.2	72.0

K10+ days out of role**Table S-50: Number of days out of role (K10+) among 11-17 year-olds by sex**

Days out of role	Males (%)	Females (%)
0 days	75.7	65.6
1 day	7.8	8.3
2 days	5.5	8.7
3 days	3.0	4.8
4 days	2.6	4.6
5-9 days	3.4	5.0
10 or more days	1.9	3.0

Self-harm

Table S-51: Self harm among 12-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)	Received medical treatment for self-harm in previous 12 months (%)
Males	Total difficulties rating—				
	Normal	4.5	1.5	2.5	np
	Borderline	10.2	5.1	6.3	np
	Abnormal	23.4	11.3	15.1	np
	Subscales in abnormal range—				
	Emotional symptoms	22.5	11.3	16.7	np
	Conduct problems	19.8	8.5	12.5	np
Hyperactivity	9.4	4.4	7.5	np	
Peer problems	32.4	12.0	19.4	np	
Females	Total difficulties rating—				
	Normal	7.2	3.1	5.0	np
	Borderline	27.7	15.0	23.6	np
	Abnormal	46.7	36.4	40.2	6.9
	Subscales in abnormal range—				
	Emotional symptoms	41.4	29.0	36.9	4.8
	Conduct problems	38.7	30.0	36.6	6.9
Hyperactivity	31.0	21.5	25.1	3.3	
Peer problems	52.2	39.2	46.4	6.9	

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-52: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and family type

Major depressive disorder status	Family type	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)
Major depressive disorder based on adolescent report	Families with two parents or carers	50.1	37.3	44.7
	Original family	46.3	36.5	40.4
	Step family	55.5	33.1	50.3
	Blended family	66.5	49.8	63.0
	Other family (a)	54.5	np	54.5
	Families with one parent or carer	62.3	42.9	50.3
No major depressive disorder based on adolescent report	Families with two parents or carers	6.4	2.5	4.2
	Original family	5.4	1.8	3.6
	Step family	14.7	7.8	8.0
	Blended family	9.2	4.8	6.3
	Other family (a)	np	np	np
	Families with one parent or carer	8.4	3.6	5.1
All persons	Families with two parents or carers	9.7	5.1	7.3
	Original family	8.0	4.0	5.9
	Step family	21.8	12.2	15.4
	Blended family	13.9	8.4	11.0
	Other family (a)	21.4	15.1	21.4
	Families with one parent or carer	14.9	8.3	10.5

'Original family' has at least one child living with their natural, adoptive or foster parents, and no step children.

'Other family' is where all children are not the natural, adopted, foster or step child of one or both carers.

(a) Data to be treated with caution due to low respondent numbers in this category.

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-53: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and household income

Major depressive disorder status	Household income before tax	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)
Major depressive disorder based on adolescent report	\$130,000 or more per year	47.5	38.9	41.9
	\$52,000-\$129,999 per year	51.4	36.1	46.2
	Less than \$52,000 per year	63.7	44.5	52.1
No major depressive disorder based on adolescent report	\$130,000 or more per year	6.1	2.8	4.3
	\$52,000-\$129,999 per year	7.3	2.6	4.5
	Less than \$52,000 per year	7.0	2.6	4.4
All persons	\$130,000 or more per year	8.8	5.2	6.7
	\$52,000-\$129,999 per year	11.0	5.5	8.0
	Less than \$52,000 per year	13.3	7.2	9.7

Household income includes the combined income for the 2011-12 financial year of everyone living in the household before tax and other deductions are taken out.

Table S-54: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer education

Major depressive disorder status	Highest level of education of either primary or secondary parent or carer	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)
Major depressive disorder based on adolescent report	Bachelor degree or higher	48.5	35.7	43.1
	Diploma or certificate III/IV	63.1	46.8	53.9
	Year 11 or 12	59.2	41.9	47.6
	Year 10 or below	36.1	21.8	31.5
No major depressive disorder based on adolescent report	Bachelor degree or higher	6.3	2.4	3.9
	Diploma or certificate III/IV	7.4	2.9	4.5
	Year 11 or 12	5.5	1.9	4.3
	Year 10 or below	8.9	4.8	6.0
All persons	Bachelor degree or higher	9.8	5.1	7.1
	Diploma or certificate III/IV	12.2	6.7	8.8
	Year 11 or 12	9.3	4.7	7.3
	Year 10 or below	12.2	6.9	9.2

Table S-55: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer labour force status

Major depressive disorder status	Parent or carer labour force status	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)
Major depressive disorder based on adolescent report	Both parents or carers employed	51.7	40.0	45.8
	One parent or carer employed, one parent or carer not in employment	44.7	33.0	37.4
	Both parents or carers not in employment	60.7	np	60.7
	Sole parent or carer employed	59.9	36.4	48.7
	Sole parent or carer not in employment	61.2	50.4	51.5
No major depressive disorder based on adolescent report	Both parents or carers employed	6.5	2.6	4.4
	One parent or carer employed, one parent or carer not in employment	6.4	2.0	4.0
	Both parents or carers not in employment	6.1	np	np
	Sole parent or carer employed	8.2	3.5	4.9
	Sole parent or carer not in employment	7.8	4.0	4.3
All persons	Both parents or carers employed	9.5	5.2	7.2
	One parent or carer employed, one parent or carer not in employment	9.3	4.3	6.4
	Both parents or carers not in employment	14.0	6.7	11.3
	Sole parent or carer employed	14.6	7.6	10.4
	Sole parent or carer not in employment	15.8	11.0	11.4

'Not in employment' combines unemployed and not in the labour force.

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-56: Self-harm among 12-17 year-olds by adolescent reported major depressive disorder status and area of residence

Major depressive disorder status	Area of residence	Self-harm ever (%)	Self-harm 4 or more times (%)	Self-harm in previous 12 months (%)
Major depressive disorder based on adolescent report	Greater capital cities	51.5	37.7	45.4
	Rest of state	58.7	41.8	48.6
No major depressive disorder based on adolescent report	Greater capital cities	6.7	2.5	4.3
	Rest of state	7.0	3.2	4.4
All persons	Greater capital cities	10.7	5.6	8.0
	Rest of state	11.2	6.3	8.1

Based on the ABS classification Greater Capital City Statistical Area (GCCSA).

Suicidal behaviours

Table S-57: Suicide ideation, suicide plans and suicide attempts among 12-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Suicidal ideation past 12 months (%)	Suicide plan past 12 months (%)	Suicide attempt in past 12 months (%)
Males	Total difficulties rating—			
	Normal	2.6	1.4	0.6
	Borderline	8.9	6.0	3.6
	Abnormal	15.5	12.6	6.4
	Subscales in abnormal range—			
	Emotional symptoms	17.0	16.3	9.7
	Conduct problems	15.3	11.1	5.1
Females	Hyperactivity	9.7	7.4	3.1
	Peer problems	16.5	14.0	7.8
	Total difficulties rating—			
	Normal	4.5	2.8	0.8
	Borderline	18.8	13.4	3.1
	Abnormal	36.3	28.1	18.3
	Subscales in abnormal range—			
Emotional symptoms	33.4	25.9	13.7	
Conduct problems	33.2	28.4	19.2	
Hyperactivity	19.5	15.6	8.0	
Peer problems	39.8	34.4	19.4	

Table S-58: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and family type

Major depressive disorder status	Family type	Suicidal ideation (%)	Suicide plan (%)	Suicide attempt (%)
Major depressive disorder based on adolescent report	Families with two parents or carers	47.0	37.4	17.4
	Original family	46.9	36.5	16.1
	Step family	36.8	34.1	np
	Blended family	68.5	54.9	np
	Other family (a)	np	np	np
	Families with one parent or carer	52.0	44.5	24.6
No major depressive disorder based on adolescent report	Families with two parents or carers	3.6	2.0	0.7
	Original family	3.5	1.9	0.7
	Step family	5.1	np	np
	Blended family	3.9	np	np
	Other family (a)	np	np	np
	Families with one parent or carer	3.6	1.7	1.0
All persons	Families with two parents or carers	6.9	4.7	2.0
	Original family	6.3	4.1	1.7
	Step family	10.7	8.0	3.4
	Blended family	9.2	6.6	3.4
	Other family (a)	12.7	np	np
	Families with one parent or carer	9.5	6.8	3.8

'Original family' has at least one child living with their natural, adoptive or foster parents, and no step children.

'Other family' is where all children are not the natural, adopted, foster or step child of one or both carers.

(a) Data to be treated with caution due to low respondent numbers in this category.

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-59: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and household income

Major depressive disorder status	Household income before tax	Suicidal ideation (%)	Suicide plan (%)	Suicide attempt (%)
Major depressive disorder based on adolescent report	\$130,000 or more per year	47.1	37.3	8.7
	\$52,000-\$129,999 per year	46.7	37.5	19.1
	Less than \$52,000 per year	56.8	47.8	28.6
No major depressive disorder based on adolescent report	\$130,000 or more per year	3.0	1.8	0.4
	\$52,000-\$129,999 per year	3.9	1.9	0.7
	Less than \$52,000 per year	4.2	2.3	1.4
All persons	\$130,000 or more per year	5.9	4.1	1.0
	\$52,000-\$129,999 per year	7.5	4.9	2.3
	Less than \$52,000 per year	10.0	7.3	4.4

Household income includes the combined income for the 2011-12 financial year of everyone living in the household before tax and other deductions are taken out.

Table S-60: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer education

Major depressive disorder status	Highest level of education of either primary or secondary parent or carer	Suicidal ideation (%)	Suicide plan (%)	Suicide attempt (%)
Major depressive disorder based on adolescent report	Bachelor degree or higher	43.6	31.8	14.3
	Diploma or certificate III/IV	54.7	47.0	20.6
	Year 11 or 12	44.0	41.5	25.8
	Year 10 or below	47.3	37.6	27.3
No major depressive disorder based on adolescent report	Bachelor degree or higher	3.3	1.9	0.7
	Diploma or certificate III/IV	3.4	1.2	0.7
	Year 11 or 12	3.8	2.9	np
	Year 10 or below	6.1	3.8	np
All persons	Bachelor degree or higher	6.6	4.4	1.8
	Diploma or certificate III/IV	7.9	5.2	2.4
	Year 11 or 12	6.6	5.7	2.3
	Year 10 or below	11.2	8.0	5.3

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-61: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and parent or carer labour force status

Major depressive disorder status	Parent or carer labour force status	Suicidal ideation (%)	Suicide plan (%)	Suicide attempt (%)
Major depressive disorder based on adolescent report	Both parents or carers employed	49.6	39.3	17.1
	One parent or carer employed, one parent or carer not in employment	35.8	29.9	12.4
	Both parents or carers not in employment	75.1	68.2	55.2
	Sole parent or carer employed	51.7	38.8	21.8
	Sole parent or carer not in employment	46.8	44.5	20.6
No major depressive disorder based on adolescent report	Both parents or carers employed	3.2	1.9	0.6
	One parent or carer employed, one parent or carer not in employment	5.2	2.1	1.3
	Both parents or carers not in employment	5.3	np	np
	Sole parent or carer employed	1.9	1.2	np
	Sole parent or carer not in employment	5.7	np	np
All persons	Both parents or carers employed	6.3	4.4	1.7
	One parent or carer employed, one parent or carer not in employment	7.5	4.1	2.1
	Both parents or carers not in employment	15.3	12.6	7.9
	Sole parent or carer employed	8.1	5.9	3.1
	Sole parent or carer not in employment	11.9	9.0	5.0

'Not in employment' combines unemployed and not in the labour force.

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-62: Suicidal ideation, suicide plans and suicide attempts among 12-17 year-olds by adolescent reported major depressive disorder status and area of residence

Major depressive disorder status	Area of residence	Suicidal ideation (%)	Suicide plan (%)	Suicide attempt (%)
Major depressive disorder based on adolescent report	Greater capital cities	46.3	34.9	16.3
	Rest of state	52.8	48.5	25.9
No major depressive disorder based on adolescent report	Greater capital cities	3.3	1.7	0.8
	Rest of state	4.2	2.3	0.8
All persons	Greater capital cities	7.1	4.7	2.2
	Rest of state	8.2	6.0	2.8

Based on the ABS classification Greater Capital City Statistical Area (GCCSA).

Smoking, alcohol, and cannabis use and sexual behaviours by Strengths and Difficulties Questionnaire scales

Table S-63: Smoking among 13-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Ever smoked at least once a week (%)	Smoked in last 30 days (%)
Males	Total difficulties rating—		
	Normal	7.2	5.2
	Borderline	14.2	9.6
	Abnormal	13.5	10.4
	Subscales in abnormal range—		
	Emotional symptoms	8.9	8.9
	Conduct problems	23.5	15.4
Females	Hyperactivity	15.5	11.3
	Peer problems	np	np
	Total difficulties rating—		
	Normal	6.0	3.5
	Borderline	12.5	10.6
	Abnormal	35.6	28.9
	Subscales in abnormal range—		
Emotional symptoms	23.3	18.9	
Conduct problems	37.1	30.1	
Hyperactivity	24.1	18.2	
Peer problems	30.0	25.9	

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-64: Alcohol consumption among 13-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Ever drunk alcohol (%)	Drunk alcohol in last 30 days (%)	More than 4 drinks in a row in last 30 days (%)
Males	Total difficulties rating—			
	Normal	35.7	17.1	11.9
	Borderline	42.7	20.0	13.9
	Abnormal	41.4	21.9	17.6
	Subscales in abnormal range—			
	Emotional symptoms	34.7	14.8	9.6
	Conduct problems	50.7	28.2	22.8
Females	Hyperactivity	45.9	24.2	16.0
	Peer problems	31.6	13.4	7.8
	Total difficulties rating—			
	Normal	33.5	14.5	9.4
	Borderline	42.5	23.7	17.6
	Abnormal	60.3	31.2	22.3
	Subscales in abnormal range—			
Emotional symptoms	53.9	27.2	20.5	
Conduct problems	63.4	34.9	25.7	
Hyperactivity	52.0	26.8	18.8	
Peer problems	51.6	23.0	14.2	

Table S-65: Cannabis and other drug use among 13-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Ever used cannabis (%)	Used cannabis in last 30 days (%)	Ever used other drugs (%)	Used other drugs in last 30 days (%)
Males	Total difficulties rating—				
	Normal	10.1	4.0	2.7	0.5
	Borderline	15.7	8.9	7.4	np
	Abnormal	25.2	14.6	9.6	4.3
	Subscales in abnormal range—				
	Emotional symptoms	18.3	7.8	7.2	np
	Conduct problems	28.1	21.2	13.6	5.0
Females	Hyperactivity	21.8	12.4	8.2	3.4
	Peer problems	12.5	np	np	np
	Total difficulties rating—				
	Normal	7.6	2.1	2.3	0.7
	Borderline	11.0	6.4	7.2	5.5
	Abnormal	28.3	13.6	17.3	5.7
	Subscales in abnormal range—				
Emotional symptoms	21.6	9.0	11.1	4.0	
Conduct problems	30.9	12.0	20.3	6.2	
Hyperactivity	20.2	9.9	14.8	7.4	
Peer problems	21.4	9.9	11.1	np	

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-66: Sexual behaviours among 13-17 year-olds by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Ever had sexual intercourse (%)	Had sexual intercourse with 4 or more persons (%)
Males	Total difficulties rating—		
	Normal	12.6	3.2
	Borderline	14.1	8.2
	Abnormal	17.5	8.3
	Subscales in abnormal range—		
	Emotional symptoms	18.2	5.7
	Conduct problems	17.5	11.0
Females	Hyperactivity	14.6	7.5
	Peer problems	13.4	np
	Total difficulties rating—		
	Normal	12.5	2.1
	Borderline	18.2	3.8
	Abnormal	35.6	14.3
	Subscales in abnormal range—		
Emotional symptoms	29.8	8.9	
Conduct problems	31.9	16.0	
Hyperactivity	24.4	10.1	
Peer problems	33.0	15.5	

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-67: Use of protection during last sexual intercourse or consumption of alcohol or drugs prior to last sexual intercourse among 13-17 year-olds who have ever had sexual intercourse by sex and adolescent self-reported SDQ rating

Sex	SDQ scale rating and subscales	Used a condom (%)	Used birth control pills (%)	Did not use any method to prevent pregnancy or sexually transmitted infections (%)	Drank alcohol or used drugs (%)
Males	Total difficulties rating—				
	Normal	76.8	31.9	7.5	21.1
	Borderline	47.1	26.5	24.7	37.8
	Abnormal	49.9	44.9	np	49.2
	Subscales in abnormal range—				
	Emotional symptoms	53.8	41.0	np	42.6
	Conduct problems	45.9	26.4	24.3	45.9
Females	Hyperactivity	51.4	30.0	16.6	43.6
	Peer problems	np	np	np	57.6
	Total difficulties rating—				
	Normal	69.3	41.9	11.1	15.4
	Borderline	64.4	54.6	np	25.4
	Abnormal	51.0	47.5	13.8	30.8
	Subscales in abnormal range—				
Emotional symptoms	50.1	47.9	15.3	25.7	
Conduct problems	47.1	34.2	23.9	32.7	
Hyperactivity	61.1	43.1	22.3	37.4	
Peer problems	35.7	38.6	19.1	20.7	

np Not available for publication because of small cell size, but included in totals where applicable.

Problem eating behaviours

Table S-68: Problem eating behaviours and weight control behaviours in the past 12 months among 11-17 year-olds by sex and age group

Age group	Eating behaviour	Males (%)	Females (%)	Persons (%)
11-15 years	Went on a diet to control weight	20.7	31.7	25.9
	Went on an eating binge	23.2	25.7	24.4
	Went on an eating binge, at least once a week	3.4	4.3	3.8
	Regularly exercised when supposed to be doing other things, or while injured, in order to control weight	35.8	39.1	37.3
	Fasted for at least a day to control weight	12.4	15.4	13.8
	Fasted for at least a day to control weight, at least once a week	4.1	5.3	4.7
	Vomited to control weight	3.4	4.5	3.9
	Vomited to control weight, at least once a week	0.7	1.2	0.9
	Took laxatives to control weight	2.9	2.7	2.8
	Took laxatives to control weight, at least once a week	0.7	1.0	0.9
	Vomited or took laxatives to control weight	5.5	6.1	5.8
	Vomited or took laxatives to control weight, at least once a week	1.3	1.8	1.5
	Low weight problem eating behaviours	0.8	1.3	1.1
	Binge eating and purging problem eating behaviours	1.0	1.1	1.0
	16-17 years	Went on a diet to control weight	15.2	45.1
Went on an eating binge		35.1	37.8	36.5
Went on an eating binge, at least once a week		6.7	10.0	8.4
Regularly exercised when supposed to be doing other things, or while injured, in order to control weight		28.4	38.9	33.8
Fasted for at least a day to control weight		9.9	24.5	17.4
Fasted for at least a day to control weight, at least once a week		1.8	7.6	4.8
Vomited to control weight		1.7	10.8	6.4
Vomited to control weight, at least once a week		np	3.0	1.9
Took laxatives to control weight		1.6	5.3	3.5
Took laxatives to control weight, at least once a week		np	1.9	1.3
Vomited or took laxatives to control weight		2.3	13.5	8.1

Age group	Eating behaviour	Males (%)	Females (%)	Persons (%)
	Vomited or took laxatives to control weight, at least once a week	1.0	4.7	2.9
	Low weight problem eating behaviours	np	2.0	1.1
	Binge eating and purging problem eating behaviours	np	3.0	1.8
11-17 years	Went on a diet to control weight	19.2	35.8	27.2
	Went on an eating binge	26.5	29.4	27.9
	Went on an eating binge, at least once a week	4.3	6.0	5.1
	Regularly exercised when supposed to be doing other things, or while injured, in order to control weight	33.7	39.0	36.3
	Fasted for at least a day to control weight	11.7	18.2	14.9
	Fasted for at least a day to control weight, at least once a week	3.5	6.0	4.7
	Vomited to control weight	3.0	6.4	4.6
	Vomited to control weight, at least once a week	0.7	1.8	1.2
	Took laxatives to control weight	2.5	3.5	3.0
	Took laxatives to control weight, at least once a week	0.7	1.3	1.0
	Vomited or took laxatives to control weight	4.6	8.4	6.5
	Vomited or took laxatives to control weight, at least once a week	1.2	2.7	1.9
	Low weight problem eating behaviours	0.7	1.5	1.1
	Binge eating and purging problem eating behaviours	0.9	1.7	1.3

np Not available for publication because of small cell size, but included in totals where applicable.

Bullying

Table S-69: Frequency of bullying in previous 12 months in 11-17 year-olds by sex and age group

Age group	Frequency of bullying	Males (%)	Females (%)	Persons (%)
11-15 years	I was not bullied in the last 12 months	65.8	60.2	63.2
	Once or twice in the last 12 months	19.9	20.3	20.1
	I was bullied every few months	4.6	6.7	5.6
	I was bullied every few weeks	3.0	4.7	3.8
	I was bullied about once a week	2.9	2.6	2.8
	I was bullied most days	3.8	5.4	4.6
16-17 years	I was not bullied in the last 12 months	75.3	68.9	72.0
	Once or twice in the last 12 months	14.4	18.7	16.6
	I was bullied every few months	3.8	4.6	4.2
	I was bullied every few weeks	2.9	2.6	2.7
	I was bullied about once a week	1.1	2.4	1.8
	I was bullied most days	2.5	2.7	2.6
11-17 years	I was not bullied in the last 12 months	68.4	62.9	65.7
	Once or twice in the last 12 months	18.4	19.8	19.1
	I was bullied every few months	4.4	6.0	5.2
	I was bullied every few weeks	3.0	4.1	3.5
	I was bullied about once a week	2.4	2.6	2.5
	I was bullied most days	3.5	4.6	4.0

Internet use and electronic games

Table S-70: Time spent using the internet or playing electronic games among 11-17 year-olds by sex and age group

Sex	Age group	Average time spent per day	Internet use on weekdays (%)	Internet use on weekends (%)	Electronic games on weekdays (%)	Electronic games on weekends (%)
Males	11-15 years	Doesn't use	1.2	1.2	3.7	3.7
		1-2 hours	52.8	41.2	64.6	40.8
		3-4 hours	23.0	27.2	16.1	27.6
		5-8 hours	13.8	20.0	11.0	19.7
		9 hours or more	9.1	10.4	4.5	8.2
	16-17 years	Doesn't use	np	np	9.4	9.4
		1-2 hours	33.4	29.1	62.6	43.2
		3-4 hours	27.9	21.6	15.2	23.0
		5-8 hours	22.0	31.8	9.8	17.5
		9 hours or more	16.1	16.9	3.0	6.9
Females	11-15 years	Doesn't use	1.5	1.5	19.0	19.0
		1-2 hours	52.5	42.6	67.1	59.7
		3-4 hours	22.6	25.5	9.9	14.6
		5-8 hours	16.5	20.9	3.0	5.4
		9 hours or more	6.9	9.5	1.0	1.3
	16-17 years	Doesn't use	np	np	37.6	37.6
		1-2 hours	27.5	22.8	54.1	47.5
		3-4 hours	32.1	27.8	4.9	8.5
		5-8 hours	25.2	31.5	2.7	4.6
		9 hours or more	15.0	17.6	np	1.8
Persons	11-15 years	Doesn't use	1.3	1.3	11.0	11.0
		1-2 hours	52.7	41.9	65.8	49.7
		3-4 hours	22.8	26.4	13.2	21.5
		5-8 hours	15.1	20.4	7.2	12.9
		9 hours or more	8.1	10.0	2.8	4.9
	16-17 years	Doesn't use	0.4	0.4	23.9	23.9
		1-2 hours	30.3	25.9	58.2	45.4
		3-4 hours	30.0	24.8	9.9	15.6
		5-8 hours	23.7	31.7	6.1	10.8
		9 hours or more	15.5	17.3	1.8	4.2

np Not available for publication because of small cell size, but included in totals where applicable.

Table S-71: Problem internet or electronic gaming behaviours among 11-17 year-olds by sex and age group

Age group	Indicators of problem behaviour related to internet use or electronic gaming	Males (%)	Females (%)	Persons (%)
11-15 years	Went without eating or sleeping	5.2	4.8	5.0
	Feel bothered when not doing	22.0	18.2	20.2
	Use when not really interested	29.0	26.3	27.7
	Spent less time than should with family or friends, doing school work or work	20.1	16.9	18.6
	Tried unsuccessfully to spend less time	25.2	24.3	24.8
	Problem internet or electronic gaming behaviours	3.5	3.0	3.3
16-17 years	Went without eating or sleeping	8.4	7.7	8.1
	Feel bothered when not doing	20.2	23.4	21.8
	Use when not really interested	39.6	44.8	42.3
	Spent less time than should with family or friends, doing school work or work	24.5	32.8	28.8
	Tried unsuccessfully to spend less time	17.6	22.7	20.2
	Problem internet or electronic gaming behaviours	4.4	6.5	5.5
11-17 years	Went without eating or sleeping	6.1	5.7	5.9
	Feel bothered when not doing	21.5	19.8	20.7
	Use when not really interested	31.9	32.0	32.0
	Spent less time than should with family or friends, doing school work or work	21.3	21.8	21.6
	Tried unsuccessfully to spend less time	23.1	23.8	23.5
	Problem internet or electronic gaming behaviours	3.8	4.1	3.9

Table S-72: Problem internet use or electronic gaming behaviours among 11-17 year-olds by sex and adolescent self-reported SDQ rating

SDQ scale rating and subscales	Males (%)	Females (%)	Persons (%)
Total difficulties rating—			
Normal	2.0	2.2	2.1
Borderline	7.3	6.6	7.0
Abnormal	14.3	12.9	13.5
Subscales in abnormal range—			
Emotional symptoms	16.8	12.2	13.4
Conduct problems	11.3	12.3	11.8
Hyperactivity	11.6	10.7	11.2
Peer problems	6.7	10.8	9.1

www.health.gov.au

All information in this publication is correct as at August 2015